

Spherical Harmonics

David Eberly
Geometric Tools, LLC
<http://www.geometrictools.com/>
Copyright © 1998-2008. All Rights Reserved.

Created: March 2, 1999
Last Modified: March 2, 2008

Contents

1 Discussion

2

1 Discussion

This came up in verifying some potential equations arising from fast solution of the n -body problem.

Associated Lagrange functions are $P_n^m(u)$ for $n \geq 0$ and $-n \leq m \leq n$. Index n is called the degree, index m is called the order. They are defined by the following.

- Zero order:

$$P_n^0(u) = \frac{(-1)^n}{2^n n!} \frac{d^n}{du^n} (1 - u^2)^n =: P_n(u)$$

- Positive order:

$$P_n^m(u) = (1 - u^2)^{m/2} \frac{d^m}{du^m} P_n(u), \quad m > 0$$

- Negative order:

$$P_n^{-m}(u) = (-1)^m \frac{(n - m)!}{(n + m)!} P_n^m(u), \quad m > 0$$

To compute the nonnegative order terms (in a memoized way), use the following scheme. The negative order terms are then easily constructed using the definition above. First note that

$$P_n^n(u) = \frac{(1 - u^2)^{n/2}}{2^n n!} \frac{d^{2n}}{du^{2n}} (u^2 - 1)^n = \frac{(2n)!(1 - u^2)^{n/2}}{2^n n!}. \quad (1)$$

for $n \geq 0$. These are the initial conditions. The recursive formulas are

$$P_n^m(u) = \frac{(2n - 1)uP_{n-1}^m(u) - (n + m - 1)P_{n-2}^m(u)}{n - m} \quad (2)$$

for $n \geq 2$, and

$$P_n^m(u) = \frac{(1 - u^2)^{1/2}}{2mu} (P_n^{m+1}(u) + (n + m)(n - m + 1)P_n^{m-1}(u)) \quad (3)$$

for $m \geq 1$. Let $P(n, m)$ represent the value of $P_n^m(u)$. The pseudocode to evaluate all of $P_N^m(u)$ for specified N and u is

```

P(0,0) = 1;
P(1,0) = u;
P(1,1) = sqrt(1-u*u);
for (n = 2; n <= N; n++)
{
    evaluate P(n,n) using equation (1);
    for (m = 0; m <= n-2; m++)
        evaluate P(n,m) using equation (2);
    evaluate P(n,n-1) using equation (3);
}

```

The following table shows values for $P_n^m(u)$ for $n \leq 3$.

$$\begin{array}{llll}
 & & & P_3^3 = 15(1-u^2)^{3/2} \\
 & & P_2^2 = 3(1-u^2) & P_3^2 = 15u(1-u^2) \\
 P_1^1 = (1-u^2)^{1/2} & P_2^1 = 3u(1-u^2)^{1/2} & P_3^1 = \frac{3}{2}(5u^2-1)(1-u^2)^{1/2} & \\
 P_0^0 = 1 & P_1^0 = u & P_2^0 = \frac{1}{2}(3u^2-1) & P_3^0 = \frac{1}{2}u(5u^2-3) \\
 P_1^{-1} = -\frac{1}{2}(1-u^2)^{1/2} & P_2^{-1} = -\frac{1}{2}u(1-u^2)^{1/2} & P_3^{-1} = -\frac{1}{8}(5u^2-1)(1-u^2)^{1/2} & \\
 & P_2^{-2} = \frac{1}{8}(1-u^2) & P_3^{-2} = \frac{1}{8}u(1-u^2) & \\
 & & P_3^{-3} = -\frac{1}{48}(1-u^2)^{3/2} &
 \end{array}$$

The derivatives of the associated Legendre functions can be computed using the following formula:

$$\frac{dP_n^m}{du} = \frac{muP_n^m - (n+m)(n-m+1)\sqrt{1-u^2}P_n^{m-1}}{1-u^2}. \quad (4)$$

When $m = 0$, the formula involves functions of order -1 . If you want to compute derivatives only using functions of nonnegative order, use the definition for the negative order functions:

$$\frac{dP_n}{du} = \frac{-n(n+1)P_n^{-1}}{\sqrt{1-u^2}} = \frac{P_n^1}{\sqrt{1-u^2}}. \quad (5)$$

Let $DP(n,m)$ represent the value of $dP_n^m(u)/du$. The pseudocode to evaluate all of $dP_N^m(u)/du$ for specified N and u is as follows. First compute all of $P_n^m(u)$ for $n \leq N$; then use

```

DP(0,0) = 0;
for (n = 1; n <= N; n++) {
    evaluate DP(n,0) using equation (5)
    for (m = 1; m <= n; m++)
        evaluate DP(n,m) using equation (4)
}

```

For derivatives of negative order functions, just use the definition relating the negative order functions to the positive order functions.

The following table shows values for $dP_n^m(u)/du$ for $n \leq 3$.

$$\begin{array}{lll}
DP_3^3 = -45(1-u^2)^{1/2} & & \\
DP_2^2 = -6u & & DP_3^2 = 15(1-3u^2) \\
DP_1^1 = \frac{-u}{\sqrt{1-u^2}} & DP_2^1 = \frac{3(1-2u^2)}{\sqrt{1-u^2}} & DP_3^1 = \frac{3}{2} \frac{u(11-15u^2)}{\sqrt{1-u^2}} \\
DP_0^0 = 0 & DP_1^0 = 1 & DP_2^0 = 3u & DP_3^0 = \frac{3}{2}(5u^2-1) \\
DP_1^{-1} = \frac{1}{2} \frac{u}{\sqrt{1-u^2}} & DP_2^{-1} = -\frac{1}{2} \frac{(1-2u^2)}{\sqrt{1-u^2}} & DP_3^{-1} = -\frac{1}{8} \frac{u(11-15u^2)}{\sqrt{1-u^2}} \\
& DP_2^{-2} = -\frac{1}{4}u & DP_3^{-2} = \frac{1}{8}(1-3u^2) \\
& & DP_3^{-3} = \frac{1}{16}u(1-u^2)^{1/2}
\end{array}$$

Note that the functions $dP_n^{\pm 1}/du$ are unbounded at $u = \pm 1$, so you may have numerical problems to deal with at those points.

The spherical harmonic functions are defined by

$$Y_n^m(\theta, \phi) = \sqrt{\frac{(n-|m|)!}{(n+|m|)!}} P_n^m(\cos \theta) \exp(im\phi)$$

for $n \geq 0$ and $-n \leq m \leq n$. Spherical coordinates are $x = r \cos \phi \sin \theta$, $y = r \sin \phi \sin \theta$, and $z = r \cos \theta$. The potential function $\Phi(r, \theta, \phi)$ which satisfies Laplaces equation

$$\frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \Phi) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial \Phi}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 \Phi}{\partial \phi^2} = 0$$

is

$$\Phi = \sum_{n=0}^{\infty} \sum_{m=-n}^n \left(L_n^m r^n + M_n^m r^{-(n+1)} \right) Y_n^m(\theta, \phi).$$

The L constants are 0 for the multipole expansion; the M constants are 0 for the local expansion.

To compute the gradient of Φ , in spherical coordinates,

$$\nabla \Phi = \frac{\partial \Phi}{\partial r} \vec{e}_r + \frac{1}{r} \frac{\partial \Phi}{\partial \theta} \vec{e}_\theta + \frac{1}{r \sin \theta} \frac{\partial \Phi}{\partial \phi} \vec{e}_\phi$$

where $\vec{e}_r = (\cos \phi \sin \theta, \sin \phi \sin \theta, \cos \theta)$, $\vec{e}_\theta = (\cos \phi \cos \theta, \sin \phi \cos \theta, -\sin \theta)$, and $\vec{e}_\phi = (-\sin \phi, \cos \phi, 0)$. The r derivative is

$$\begin{aligned}
\frac{\partial \Phi}{\partial r} &= \sum_{n=0}^{\infty} \sum_{m=-n}^n (n L_n^m r^{n-1} - (n+1) M_n^m r^{-(n+2)}) Y_n^m(\theta, \phi) \\
&= \sum_{n=0}^{\infty} \sum_{m=-n}^n (n L_n^m r^{n-1} - (n+1) M_n^m r^{-(n+2)}) \sqrt{\frac{(n-|m|)!}{(n+|m|)!}} P_n^m(\cos \theta) \exp(im\phi).
\end{aligned}$$

You need to compute the functions $P_n^m(\cos \theta)$ using the recursions given earlier, where the evaluation point is $u = \cos \theta$. The ϕ derivative is

$$\begin{aligned}
\frac{\partial \Phi}{\partial \phi} &= \sum_{n=0}^{\infty} \sum_{m=-n}^n (L_n^m r^n + M_n^m r^{-(n+1)}) \frac{\partial Y_n^m(\theta, \phi)}{\partial \phi} \\
&= \sum_{n=0}^{\infty} \sum_{m=-n}^n (L_n^m r^n + M_n^m r^{-(n+1)}) im \sqrt{\frac{(n-|m|)!}{(n+|m|)!}} P_n^m(\cos \theta) \exp(im\phi).
\end{aligned}$$

Again, you need only compute the functions $P_n^m(\cos \theta)$ using the recursions with $u = \cos \theta$. The θ derivative is

$$\begin{aligned} \frac{\partial \Phi}{\partial \theta} &= \sum_{n=0}^{\infty} \sum_{m=-n}^n (L_n^m r^n + M_n^m r^{-(n+1)}) \frac{\partial Y_n^m(\theta, \phi)}{\partial \theta} \\ &= \sum_{n=0}^{\infty} \sum_{m=-n}^n (L_n^m r^n + M_n^m r^{-(n+1)}) \sqrt{\frac{(n-|m|)!}{(n+|m|)!}} \frac{dP_n^m(\cos \theta)}{du} (-\sin \theta) \exp(im\phi). \end{aligned}$$

Now you need to compute the functions dP_n^m/du using the recursion formula given earlier for the derivatives of associated Legendre functions.