

Distance Between Two Circles in 3D

David Eberly

Geometric Tools, LLC

<http://www.geometrictools.com/>

Copyright © 1998-2008. All Rights Reserved.

Created: March 2, 1999

Last Modified: March 1, 2008

Contents

1 Discussion

2

1 Discussion

A circle in 3D is represented by a center \mathbf{C} , a radius R , and a plane containing the circle, $\mathbf{N} \cdot (\mathbf{X} - \mathbf{C}) = 0$ where \mathbf{N} is a unit length normal to the plane. If \mathbf{U} and \mathbf{V} are also unit length vectors so that \mathbf{U} , \mathbf{V} , and \mathbf{N} form a right-handed orthonormal coordinate system (the matrix with these vectors as columns is orthonormal with determinant 1), then the circle is parameterized as

$$\mathbf{X} = \mathbf{C} + R(\cos(\theta)\mathbf{U} + \sin(\theta)\mathbf{V}) =: \mathbf{C} + R\mathbf{W}(\theta)$$

for angles $\theta \in [0, 2\pi)$. Note that $|\mathbf{X} - \mathbf{C}| = R$, so the \mathbf{X} values are all equidistant from \mathbf{C} . Moreover, $\mathbf{N} \cdot (\mathbf{X} - \mathbf{C}) = 0$ since \mathbf{U} and \mathbf{V} are perpendicular to \mathbf{N} , so the \mathbf{X} lie in the plane.

Let the two circles be $\mathbf{C}_0 + R_0\mathbf{W}_0(\theta)$ for $\theta \in [0, 2\pi)$ and $\mathbf{C}_1 + R_1\mathbf{W}_1(\phi)$ for $\phi \in [0, 2\pi)$. The squared distance between any two points on the circles is

$$\begin{aligned} F(\theta, \phi) &= |\mathbf{C}_1 - \mathbf{C}_0 + R_1\mathbf{W}_1 - R_0\mathbf{W}_0|^2 \\ &= |\mathbf{D}|^2 + R_0^2 + R_1^2 + 2R_1\mathbf{D} \cdot \mathbf{W}_1 - 2R_0R_1\mathbf{W}_0 \cdot \mathbf{W}_1 - 2R_0\mathbf{D} \cdot \mathbf{W}_0 \end{aligned}$$

where $\mathbf{D} = \mathbf{C}_1 - \mathbf{C}_0$. Since F is doubly periodic and continuously differentiable, its global minimum must occur when $\nabla F = (0, 0)$. The partial derivatives are

$$\frac{\partial F}{\partial \theta} = -2R_0\mathbf{D} \cdot \mathbf{W}'_0 - 2R_0R_1\mathbf{W}'_0 \cdot \mathbf{W}_1$$

and

$$\frac{\partial F}{\partial \phi} = 2R_1\mathbf{D} \cdot \mathbf{W}'_1 - 2R_0R_1\mathbf{W}_0 \cdot \mathbf{W}'_1.$$

Define $c_0 = \cos(\theta)$, $s_0 = \sin(\theta)$, $c_1 = \cos(\phi)$, and $s_1 = \sin(\phi)$. Then $\mathbf{W}_0 = c_0\mathbf{U}_0 + s_0\mathbf{V}_0$, $\mathbf{W}_1 = c_1\mathbf{U}_1 + s_1\mathbf{V}_1$, $\mathbf{W}'_0 = -s_0\mathbf{U}_0 + c_0\mathbf{V}_0$, and $\mathbf{W}'_1 = -s_1\mathbf{U}_1 + c_1\mathbf{V}_1$. Setting the partial derivatives equal to zero leads to

$$\begin{aligned} 0 &= s_0(a_0 + a_1c_1 + a_2s_1) + c_0(a_3 + a_4c_1 + a_5s_1) \\ 0 &= s_1(b_0 + b_1c_0 + b_2s_0) + c_1(b_3 + b_4c_0 + b_5s_0) \end{aligned}$$

where

$$\begin{aligned} a_0 &= -\mathbf{D} \cdot \mathbf{U}_0, & a_1 &= -R_1\mathbf{U}_0 \cdot \mathbf{U}_1, & a_2 &= -R_1\mathbf{U}_0 \cdot \mathbf{V}_1, & a_3 &= \mathbf{D} \cdot \mathbf{V}_0, & a_4 &= R_1\mathbf{U}_1 \cdot \mathbf{V}_0, & a_5 &= R_1\mathbf{V}_0 \cdot \mathbf{V}_1, \\ b_0 &= -\mathbf{D} \cdot \mathbf{U}_1, & b_1 &= R_0\mathbf{U}_0 \cdot \mathbf{U}_1, & b_2 &= R_0\mathbf{U}_1 \cdot \mathbf{V}_0, & b_3 &= \mathbf{D} \cdot \mathbf{V}_1, & b_4 &= -R_0\mathbf{U}_0 \cdot \mathbf{V}_1, & b_5 &= -R_0\mathbf{V}_0 \cdot \mathbf{V}_1. \end{aligned}$$

In matrix form we have

$$\begin{bmatrix} m_{00} & m_{01} \\ m_{10} & m_{11} \end{bmatrix} \begin{bmatrix} s_0 \\ c_0 \end{bmatrix} = \begin{bmatrix} a_0 + a_1c_1 + a_2s_1 & a_3 + a_4c_1 + a_5s_1 \\ b_2s_1 + b_5c_1 & b_1s_1 + b_4c_1 \end{bmatrix} \begin{bmatrix} s_0 \\ c_0 \end{bmatrix} = \begin{bmatrix} 0 \\ -(b_0s_1 + b_3c_1) \end{bmatrix} = \begin{bmatrix} 0 \\ \lambda \end{bmatrix}$$

Let M denote the 2×2 matrix on the right-hand side of the equation. Multiplying by the adjoint of M yields

$$\det(M) \begin{bmatrix} s_0 \\ c_0 \end{bmatrix} = \begin{bmatrix} m_{11} & -m_{01} \\ -m_{10} & m_{00} \end{bmatrix} \begin{bmatrix} 0 \\ \lambda \end{bmatrix} = \begin{bmatrix} -m_{01}\lambda \\ m_{00}\lambda \end{bmatrix}. \quad (1)$$

Summing the squares of the vector components and using $s_0^2 + c_0^2 = 1$ yields

$$(m_{00}m_{11} - m_{01}m_{10})^2 = \lambda^2 (m_{00}^2 + m_{01}^2).$$

The above equation can be reduced to a polynomial of degree 8 whose roots $c_1 \in [-1, 1]$ are the candidates to provide the global minimum of F . Formally computing the determinant and using $s_1^2 = 1 - c_1^2$ leads to

$$m_{00}m_{11} - m_{01}m_{10} = p_0(c_1) + s_1p_1(c_1)$$

where $p_0(z) = \sum_{i=0}^2 p_{0i}z^i$ and $p_1(z) = \sum_{i=0}^1 p_{1i}z$. The coefficients are

$$\begin{aligned} p_{00} &= a_2b_1 - a_5b_2, \\ p_{01} &= a_0b_4 - a_3b_5, \\ p_{02} &= a_5b_2 - a_2b_1 + a_1b_4 - a_4b_5, \\ p_{10} &= a_0b_1 - a_3b_2, \\ p_{11} &= a_1b_1 - a_5b_5 + a_2b_4 - a_4b_2. \end{aligned}$$

Similarly,

$$m_{00}^2 + m_{01}^2 = q_0(c_1) + s_1q_1(c_1)$$

where $q_0(z) = \sum_{i=0}^2 q_{0i}z^i$ and $q_1(z) = \sum_{i=0}^1 q_{1i}z$. The coefficients are

$$\begin{aligned} q_{00} &= a_0^2 + a_2^2 + a_3^2 + a_5^2, \\ q_{01} &= 2(a_0a_1 + a_3a_4), \\ q_{02} &= a_1^2 - a_2^2 + a_4^2 - a_5^2, \\ q_{10} &= 2(a_0a_2 + a_3a_5), \\ q_{11} &= 2(a_1a_2 + a_4a_5). \end{aligned}$$

Finally,

$$\lambda^2 = r_0(c_1) + s_1r_1(c_1)$$

where $r_0(z) = \sum_{i=0}^2 r_{0i}z^i$ and $r_1(z) = \sum_{i=0}^1 r_{1i}z$. The coefficients are

$$\begin{aligned} r_{00} &= b_0^2, \\ r_{01} &= 0, \\ r_{02} &= b_3^2 - b_0^2, \\ r_{10} &= 0, \\ r_{11} &= 2b_0b_3. \end{aligned}$$

Combining these yields

$$0 = [(p_0^2 - r_0q_0) + (1 - c_1^2)(p_1^2 - r_1q_1)] + s_1 [2p_0p_1 - r_0q_1 - r_1q_0] = g_0(c_1) + s_1g_1(c_1) \quad (2)$$

where $g_0(z) = \sum_{i=0}^4 g_{0i} z^i$ and $g_1(z) = \sum_{i=0}^3 g_{1i} z^i$. The coefficients are

$$\begin{aligned}
g_{00} &= p_{00}^2 + p_{10}^2 - q_{00}r_{00} \\
g_{01} &= 2(p_{00}p_{01} + p_{10}p_{11}) - q_{01}r_{00} - q_{10}r_{11} \\
g_{02} &= p_{01}^2 + 2p_{00}p_{02} + p_{11}^2 - p_{10}^2 - q_{02}r_{00} - q_{00}r_{02} - q_{11}r_{11} \\
g_{03} &= 2(p_{01}p_{02} - p_{10}p_{11}) - q_{01}r_{02} + q_{10}r_{11} \\
g_{04} &= p_{02}^2 - p_{11}^2 - q_{02}r_{02} + q_{11}r_{11} \\
g_{10} &= 2p_{00}p_{10} - q_{10}r_{00} \\
g_{11} &= 2(p_{01}p_{10} + p_{00}p_{11}) - q_{11}r_{00} - q_{00}r_{11} \\
g_{12} &= 2(p_{02}p_{10} + p_{01}p_{11}) - q_{10}r_{02} - q_{01}r_{11} \\
g_{13} &= 2p_{02}p_{11} - q_{11}r_{02} - q_{02}r_{11}
\end{aligned}$$

We can eliminate the s_1 term by solving $g_0 = -s_1 g_1$ and squaring to obtain

$$0 = g_0^2 - (1 - c_1^2)g_1^2 = h(c_1)$$

where $h(z) = \sum_{i=0}^8 h_i z^i$. The coefficients are

$$\begin{aligned}
h_0 &= g_{00}^2 - g_{10}^2, \\
h_1 &= 2(g_{00}g_{01} - g_{10}g_{11}), \\
h_2 &= g_{01}^2 + g_{10}^2 - g_{11}^2 + 2(g_{00}g_{02} - g_{10}g_{12}), \\
h_3 &= 2(g_{01}g_{02} + g_{00}g_{03} + g_{10}g_{11} - g_{11}g_{12} - g_{10}g_{13}), \\
h_4 &= g_{02}^2 + g_{11}^2 - g_{12}^2 + 2(g_{01}g_{03} + g_{00}g_{04} + g_{10}g_{12} - g_{11}g_{13}), \\
h_5 &= 2(g_{02}g_{03} + g_{01}g_{04} + g_{11}g_{12} + g_{10}g_{13} - g_{12}g_{13}), \\
h_6 &= g_{03}^2 + g_{12}^2 - g_{13}^2 + 2(g_{02}g_{04} + g_{11}g_{13}), \\
h_7 &= 2(g_{03}g_{04} + g_{12}g_{13}), \\
h_8 &= g_{04}^2 + g_{13}^2.
\end{aligned}$$

To find the minimum squared distance, compute all the real-valued roots of $h(c_1) = 0$. For each c_1 , compute $s_1 = \pm\sqrt{1 - c_1^2}$ and choose either (or both) of these that satisfy equation (2). For each pair (c_1, s_1) , solve for (c_0, s_0) in equation (1). The main numerical issue to deal with is how close to zero is $\det(M)$. (TO DO: Show that this case only occurs when circles are parallel and \mathbf{D} is normal to both planes?)